



(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
18.12.2002 Bulletin 2002/51

(21) Application number: 02380119.4

(22) Date of filing: 11.06.2002

(51) Int Cl.7: A01N 59/00, A01N 37/36,
A01N 25/30, A61L 2/18,
A01N 59/16
// (A01N59/00, 37:36, 25:30,
59:16)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 15.06.2001 ES 200101398

(71) Applicant: Oftral, S.L.
28760 Tres Cantos (ES)

(72) Inventor: Herruzo Cabrera, Rafael
28760 Tres Cantos (Madrid) (ES)

(74) Representative: Elizaburu, Alberto de
Elizaburu S.A.
Miguel Angel, 21
28010 Madrid (ES)


(54) New desinfectant and antiseptic composition

(57) Wide spectrum disinfecting and antiseptic composition for use in the fields of human medicine, veterinary science and industry, characterised because it includes:

Hydrogen peroxide, lactic acid and halogen salts (Br, I) and/or salts of heavy metals (for example, silver halides) with surfactant agents, either cationic, like chlorhexidine and/or quaternary ammonium salts, like didecyl-methyl-polyoxy-ethyl-ammonium


propionate, chlorides of ammonium or compounds of ammonium propylamide or anionic, like lauryl sulphate, dodecyl sulphate or alkyl succinic salts, with suitable excipients, some of which may be ethyl or isopropyl alcohol, chlorhexidine, non-chlorinated quaternary ammonium salts, like didecyl-methyl-polyoxy-ethyl-ammonium propionate, combined or not with iodine, and/or its salts, together with excipients, some of which may be ethyl or isopropyl alcohol.

Figure 1: Assessment of the synergy of "Example a" 1/6 with respect to its basic components (at the same dilution), by means of the determination of the sporicidal effect over a commercial B subtilis germ-carrier



EP 1 266 571 A1

Figure 2 : Assessment of the synergy of "example b" diluted 1/4 with respect to its basic components (at the same dilution), by means of the determination of the sporicidal effect over a commercial B subtilis germ-carrier.



Description**FIELD OF THE INVENTION**

- 5 **[0001]** This invention refers to a new disinfectant and antiseptic composition.
[0002] The disinfecting component of this wide spectrum composition comprises oxidising agents, like hydrogen peroxide, lactic acid and halogen salts (Br, I) and/or heavy metals, plus surfactants, that is:
- 10 a) cationic surfactants, like quaternary ammonium and/or biguanidines, like chlorhexidine, or
 b) anionic surfactants, like lauryl sulphate, alkyl succinic or dodecyl sulphate salts.
- [0003]** All of them with the suitable excipients (alcohol, water, etc.).
[0004] The antiseptic component of the composition binds chlorhexidine to non-chlorinated quaternary ammonium like didecylmethylpolyoxyethylammonium propionate with iodine or iodinated salts and suitable excipients, like dyes, emollients and alcohol.
- 15 **[0005]** This disinfectant and antiseptic composition may be used in the field of medicine, veterinary science or industry.

BACKGROUND OF THE INVENTION

- 20 **[0006]** Disinfecting instruments for hospital use involves a key factor for the control of hospital infections. There are several methods and products used to perform said disinfecting. A review of the main methods and disinfecting products for hospital use, together with their indications and risks may be found in the tenth edition of the book by Piedrola Gil, "Medicina Preventiva y Salud Pública" (Preventive Medicine and Public Health), Ed Masson (Barcelona), in Chapter
- 25 37, pages 413-424, written by Herruzo Cabrera et al.
- [0007]** This chapter shows the advantages and drawbacks of 2% glutaraldehyde, the reference disinfectant and the need of finding a suitable alternative for it, since its toxicity makes it a problem for the users and even for patients, if the disinfected material is not thoroughly rinsed. An alternative may be the combination of lactic acid to hydrogen peroxide, since it combines non-toxicity to a wide microbicidal capacity. This was the object of patent PCT/ES98/00151, in whose claims includes 3-6% hydrogen peroxide, 5-15% lactic acid and combined with the previous mixture, a compound selected from 0.4-5% chlorhexidine or 0.1-3% lauryl sulphate of an alkaline metal and water as a balance, where all the percentages are with respect to the total composition. However, this compound is very acid, which may involve a problem for its use, and if its components are reduced by dilution, its disinfecting power is lost, so that it is necessary to reinforce this activity and reduce its acidity.
- 30 **[0008]** The present invention corrects these problems by using the similar products with ranges which are different to those claimed and adds others which are vital to considerably enhance its microbicidal effectiveness, besides providing the product with a considerable dilution capacity, cheapening it and even permitting its use as an antiseptic, in specific formulations, designed for such a purpose.

DETAILED DESCRIPTION OF THE INVENTION

- 40 **[0009]** This invention provides a disinfecting and antiseptic composition, comprising oxidising agents, like hydrogen peroxide, lactic acid and halogen salts (and/or heavy metal salts), together with surfactants, that is, a biguanidine like chlorhexidine with or without quaternary ammonium, or anionic surfactants, like lauryl sulphate, alkyl succinic or dodecyl sulphate salts. Additionally, the composition of the invention may contain water and some suitable excipients, like alcohol.
- 45 **[0010]** On the other hand, the composition is also antiseptic, based on chlorhexidine and non-chlorinated quaternary ammonium like didecyl methyl-polyoxy-ethyl-ammonium propionate, with iodine (metallic or in salts) and suitable excipients, like emollients and alcohol.
- 50 **[0011]** Hydrogen peroxide is an oxidising disinfectant, of medium activity against a plurality of micro-organisms. In its undiluted formulation, the disinfectant composition of the invention may contain 0.16-2.9% or 6.1-22.5% hydrogen peroxide with respect to the total.
- [0012]** Lactic acid, increasing the effectiveness of hydrogen peroxide may be available in an amount comprised between 0.1% and 24%.
- 55 **[0013]** Chlorhexidine is a biguanidine with a powerful antimicrobial activity, but has little effectiveness against virus and mycobacteria, such that in disinfecting it should be combined with other products. In the present invention, in the undiluted formula, it may be present at a concentration comprised between 5.1% and 18%.
- [0014]** Lauryl sulphate is an anionic surfactant with a wide detergent effect, but with reduced microbicidal power,

frequently used in soaps for washing hands. This disinfectant composition may contain it in a ratio comprised between 3.1% and 20%.

[0015] It may be substituted by another anionic agent, like alkyl succinic or dodecyl sulphate, with proportions similar to the anionic surfactants, with respect to the total, although it is preferable to reduce them to 1-12%.

5 **[0016]** The quaternary ammonium salts are cationic surfactants frequently used in disinfecting and antiseptics, although problems have been described related to their use due to bacteria becoming resistant to them, but may be very useful when combined with other products, since they aid in the destruction of bacteria walls. In its undiluted formulation, these products will be in a wide ratio, from 0.2% to 50%.

10 **[0017]** The halogens, like Br and I, are also frequently used in antiseptics and disinfecting and usually act by generating oxidising by-products of the microbial components. In the composition of this invention, they are incorporated at a ratio comprised between 0.001% and 5 of the undiluted formula.

[0018] The heavy metal salts, like silver, cerium, etc., are widely used in disinfectants and antiseptics, acting by accumulation of the latter in the bacterial cytoplasm. In the composition of this invention, they were incorporated in the range from 0.0002% to 2% in undiluted formulas.

15 **[0019]** The disinfecting and antiseptic composition of the invention may also contain a balance of water and excipients, normally used in disinfectant composition formulations, for example, food preserving agents, dyes, enzymes and alcohol.

[0020] In one of the preferred embodiments of the invention, the ranges of the basic substances are as follows:

- 20
- hydrogen peroxide 0.16-2.9% or 6.1-22.5%
 - lactic acid: 0.2-20%
 - chlorhexidine 5.1-18%
 - excipients and water (balance)

25 where all the percentages are by weight with respect to the total.

[0021] In another of the preferred embodiments of the invention, the disinfectant composition ranges are:

- 30
- hydrogen peroxide: 0.16-2.9% or 6.1-22.5%
 - lactic acid: 0.2-20%
 - alkaline metal lauryl sulphate: 3.1-20%
 - excipients and water (balance)

where all the percentages are by weight with respect to the total.

35 **[0022]** Chlorhexidine or lauryl sulphate may also be replaced by quaternary ammonium salts, or by mixing chlorhexidine with the quaternary ammonium salts to give the surfactant compound which binds to the hydrogen peroxide base, lactic acid and heavy metal salts and/or halogen salts (Br, I). Alcohol may also be added to improve evaporation and the antimicrobial effectiveness. They may also bind to enzymes like trypsin, amylase and lipase to give enzymatic disinfectants.

[0023] Another example to be used in antiseptics is the following one in an undiluted formulation:

- 40
- chlorhexidine: 0.2-18%
 - didecyl-methyl-polyoxy-ethyl-ammonium propionate: 1-45%
 - iodine (metallic or its salts): 0.01-5%
 - excipients (one of them may be ethyl or isopropyl alcohol)
- 45

[0024] The disinfecting and antiseptic composition of the invention is very useful as a wide spectrum disinfectant and especially destined to be used in any field, veterinary, industrial or human medicine, since it is very useful as a disinfecting agent for instruments (for example, high level disinfecting in certain semi-critical instruments, like flexible endoscopes) surfaces, conduits, etc.

50 **[0025]** For disinfecting, a suitable amount of the product of the invention should be applied, diluted according to the indication and quickness required over the material or surface to be disinfected. This application may be performed by any conventional method like immersion, spraying, injection, impregnation with the aid of a suitable applicator for the disinfecting composition over the conduits, surface or instruments to be disinfected, both in the field of human medicine and that of veterinary medicine or industry.

55 **[0026]** Some of its components are originally separated and combined at the time of use or dilution in water. At this time, the mixture is considered as activated and has a variable shelf-life, which in general exceeds two weeks (which may be increased with suitable preserving agents), although if contaminated objects are introduced, etc. it is not convenient to let the solution to age, making it recommendable to eliminate the dilutions at the end of the day.

[0027] On the other hand, the composition object of the invention may also be used as an antiseptic, both in human and veterinary medicine, it being possible to deposit it over the skin like other antiseptics, by immersion, application, spraying, etc. The antiseptic composition is already reconstituted (it is not necessary to mix its components "in situ") and has a minimum shelf-life of two years.

5 [0028] The disinfecting composition of the invention has a much greater synergic effect than that expected from the simple addition of the products present in it:

- In this sense, it was surprising to verify that the combination of hydrogen peroxide with lactic acid and surfactants (whether anionic or cationic), plus excipients, has a synergic effect, as may be seen in figures 1 and 2 (said synergy being marked by the arrows descending from the best of the basic components to the mixture) as well as in examples 1-9 which show that said combinations increase its microbicidal capacity and decreases the oxidising effect of hydrogen peroxide, which does not alter the instruments, hence being suitable for disinfecting any material, including hospital material, preventing the problems described in disinfecting with hydrogen peroxide in the equipment and persons using it for the disinfecting of endoscopes, haemodialysers, etc.

15 [0029] On the other hand, the antiseptic variant of this invention is also based on a synergic effect, as will be shown afterwards, due to the combination of chlorhexidine, certain quaternary ammonium salts and iodine.

[0030] Besides, if this mixture of halogens with chlorhexidine and quaternary ammonium salts, is added at a ratio of 10-30% to 60° ethyl or isopropyl alcohol (this alcohol completing the remaining solution by 90-70%), a fast action antiseptic may be obtained which effectively acts over healthy skin, eliminating the acquired flora and reducing the autochthonous one, as will be shown later.

[0031] The tests described below show that the disinfecting composition of the invention is very useful for disinfecting hospital or non-hospital material, even if complex, with rough surfaces, etc. where other disinfecting products may fail. The variants with antiseptic action are harmless for the skin, such that they may constitute good antiseptic products, improving the effectiveness and persistence of reference antiseptic action, 10% povidone-iodine.

25 [0032] The disinfecting and antiseptic composition of the invention is obtained by mixing its basic compounds in suitable amounts, stirring until total homogenisation, although in certain formulas, some of the basic compounds may only bind to the rest at the time of activation, which is usually the same time as dilution in water.

[0033] Structured in two parts, according to the main indication as antiseptics or disinfectants, the following examples are specific and illustrative embodiments of the composition of this invention and must not be considered as limiting its scope.

PART 1: DISINFECTANTS (main activity)

35 EXAMPLE 1

Preparation of a disinfecting composition

40 [0034] A disinfecting composition was prepared including:

- hydrogen peroxide: 16%
- lactic acid. 24%
- alkaline metal lauryl sulphate 20%
- halogen salts (Br, I): 0.4%

45 [0035] The composition was obtained by mixing the appropriate amounts of each component and stirring until complete homogenisation.

[0036] The efficacy was proved by carrying out the following tests:

[0037] Study of the bactericidal effect of the undiluted composition as well as when diluted to ½ and ¼.

50 [0038] Determination of the conservation of the concentrated and diluted solution activity during 2 weeks.

[0039] Instrument deterioration test.

[0040] For the determination of these three effects, the methods described by Herruzo et al in 1999 were used (Rev Stomatol Chir maxillofac 1999; 100: 299-305), which may be resumed as follows:

55 a) Bactericidal effect on instruments difficult to disinfect: No. 25 endodontic files.

These files are contaminated in a micro-organism for 30 minutes and after drying for 15 minutes in an oblique position, they are introduced in the disinfectant for 5, 10, 15, 20, etc. minutes. Afterwards, they are removed and submerged in a tube with glass beads and a disinfectant inhibitor (broth plus 6% Tween 80, plus 0.5% sodium

bisulphite and 0.5% sodium thiosulphate). The tubes are then stirred for 1 minute to detach the surviving micro-organisms, and samples are cultured over the entire surface of agar plates with the culture medium suitable for the tested micro-organism (Müller-Hinton or Saboureaud-dextrose for yeasts or HIB for mycobacteria, etc.). The control is similar in the whole test (disinfectant), except that the contaminated endodontic file is introduced in sterile distilled water and the culture is performed in progressive dilutions to count the cfu (colony forming units).

b) After preparation, the solution is left at room temperature for 7-14 days and the trial is repeated, as just described, but only with micro-organisms resistant to disinfecting: *P. Aeruginosa* and *Mycobacterium fortuitum*.

c) The effect on the metallic material is obtained by submerging a scalpel, after being removed from its sheath, in the different dilutions of the disinfectant composition. It is left at room temperature and the blackening or deterioration is observed during one week (equivalent to more than 2000 rinses of 10 minutes). It is recorded if there was any deterioration or not and the time at which it occurred. This test may be complicated if the first step consists in submerging the scalpel in blood (coming from a blood bank), since some of the disinfectants help to detach this organic material while others, fix it and collaborate in the deterioration of the scalpel.

[0041] The results are the following on 5 strains of each one of the species indicated: *S. aureus*, *S. epidermidis*, *E. faecalis*, *E. coli*, *K. pneumoniae*, *E. cloacae*, *P. mirabilis*, *Acinetobacter baumannii*, *P. aeruginosa* and *C. albicans*. Also, two *Mycobacterium fortuitum* strains were used (collections ATCC 609 and ATCC 6841). The tests were repeated five times with each one of the collection strains.

[0042] The time was ten minutes contact with the disinfectant for all cases.

Micro-organism	control cfu (mean)	test cfu (mean) (dilution 1/1)
<i>S. aureus</i>	135000	0
<i>S. epidermidis</i>	67000	0
<i>E. faecalis</i>	25000	0
<i>E. coli</i>	236000	0
<i>K. pneumoniae</i>	35000	0
<i>E. cloacae</i>	41000	0
<i>P. mirabilis</i>	39000	0
<i>A. baumannii</i>	64000	0
<i>P. aeruginosa</i>	190000	0
<i>C. albicans</i>	12500	0
<i>M. fortuitum</i> 609	78000	0
<i>M. fortuitum</i> 6841	56000	0

[0043] Similar results were obtained in 1/2 and 1/4 dilutions.

[0044] The repetition of the trial with *P. aeruginosa* and *M. fortuitum* 609 at 7 and 14 days, produced identical results, involving a stability of at least 14 days without use.

[0045] The deterioration of metallic instruments as labile as the scalpel blade, impregnated or not with organic material, was nil, indicating good cleaning capacity and harmlessness before this model.

[0046] On studying 2% glutaraldehyde for 20 minutes in parallel (which is the reference disinfectant, although toxic and for this reason is replaced by formulas not having this negative feature), similar results were obtained in all strains except in *Mycobacterium*, against which, it did not completely destroyed the inoculum (not even in the time of use, 20 minutes, that is, twice than used for the disinfectant of the invention). Moreover, when the scalpel model is studied with this alkylising agent, the influence of the need of a good rinse may be observed, since if the scalpel did not have any organic material, several days would be required for deterioration of the model, but if it were impregnated with blood, it would deteriorate in only a few hours.

EXAMPLE 2

Preparation of a disinfecting composition

[0047] A disinfecting composition containing the following was prepared:

- hydrogen peroxide 11%

- lactic acid 16%
- didecyl-methyl-polyoxy-ethyl-ammonium propionate:32%
- silver halides 0.01%

- 5 [0048] The composition was prepared by mixing suitable amounts of each components and stirring until complete homogenisation.
- [0049] The effectiveness was shown by performing the following tests:
- [0050] Study of the bactericidal effect of the undiluted composition, as well as that diluted to $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{6}$ and $\frac{1}{8}$.
- [0051] Determination of the preservation of the concentrated and diluted solution activity, during 2 weeks.
- 10 [0052] Instrumental deterioration test.
- [0053] The methodology is similar to that described in the previous example, except for the fact that the studies were performed in a time of 5 minutes instead of 10 and a greater dilution was reached, $\frac{1}{8}$.
- [0054] The results were similar to the previous ones, 0 survivors in all cases, besides remaining active for at least 2 weeks and the scalpel not deteriorating, with or without blood.
- 15 [0055] In this case, another more demanding test was used, since the sporicidal effect was assessed. It is similar to the bactericidal effect, but instead of contaminated instruments, a commercial, cellulose germ holder was used, impregnated in spores. In fact, this is the biological control used in sterilisation by ethylene oxide: spores of *B. subtilis*, at a concentration guaranteed by the manufacturer (3M) and which exceed a million spores per germ holder. The rest was similar to that described in the endodontic file method.
- 20 [0056] The results were excellent, since 5 decimal logarithms were destroyed in only 5 minutes in the $\frac{1}{1}$ to $\frac{1}{8}$ solutions. When lower dilutions (only with this germ) were also used, namely $\frac{1}{20}$ and $\frac{1}{50}$, during 10 minutes, the effect was lower, 2.6 log10 and 1.9 log10 of reduction, respectively, but it was still sufficiently useful since if we compare it with the reference disinfectant, 2% glutaraldehyde, in this time (10 minutes) it only reduced the inoculum by 1.5 log10.
- 25 [0057] In the successive examples, the number of micro-organisms used was reduced, since only 3 strains of the following species (isolated from ICU patients) were used: *P. aeruginosa*, *S. aureus*, *E. coli*, *E. faecalis*, together with the *M. fortuitum* strains of the ATCC 609 collection and *B. subtilis* spores.

EXAMPLE 3

- 30 [0058] A disinfectant composition containing the following was prepared:
- hydrogen peroxide 9%
 - lactic acid 4.9%
 - chlorhexidine 18%
 - 35 - silver halides 0.1%
- The composition was obtained by mixing the suitable amounts of each compound and stirring until complete homogenisation.
- [0059] The effectiveness was shown by performing the following tests:
- 40 [0060] Study of the bactericidal effect of the undiluted composition, as well as dilutions at $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{6}$ and $\frac{1}{8}$.
- [0061] Determination of the preservation of the concentrated and diluted solution activity, during 2 weeks.
- [0062] Instrumental deterioration test.
- [0063] The methodology is similar to that described in the previous example 2, except that the studies were performed until a lesser dilution, $\frac{1}{6}$, in 5 minutes and until $\frac{1}{8}$, in 10 minutes.
- 45 [0064] The results were similar to the previous ones, 0 survivors in all cases (plus 5 log reduction, including the spores), remaining active at least 2 weeks and not deteriorating the scalpel, with or without blood.

EXAMPLE 4

- 50 [0065] A disinfecting composition was prepared containing:
- hydrogen peroxide: 21%
 - lactic acid 18%
 - didecyl-methyl-polyoxy-ethyl-ammonium propionate 6%
 - 55 - silver halides: 0.001%
- [0066] The composition was obtained by mixing suitable amounts of each compound and stirring until complete homogenisation.

- [0067] The effectiveness was shown by performing the following tests:
 [0068] Study of the microbicidal effect of the undiluted composition, as well as 1/2, 1/4 and 1/6 dilutions.
 [0069] Determination of the preservation of the concentrated and diluted solution activity during 2 weeks.
 [0070] Instrument deterioration test.
 5 [0071] The methodology and results were similar to those of the previous example, but the 1/8 dilution was not used.

EXAMPLE 5

- [0072] A disinfecting composition was prepared, containing:

- 10
 - hydrogen peroxide: 11%
 - lactic acid: 16%
 - didecyl-methyl-polyoxy-ethyl-ammonium propionate 24%
 - chlorhexidine 12%
 15 - silver halides 0.05%

[0073] The composition was obtained by mixing suitable amounts of each compound and stirring until complete homogenisation.

- [0074] The effectiveness was shown by performing the following tests:
 20 [0075] Study of the bactericidal effect of the undiluted composition, as well as 1/2, 1/4, 1/6 and 1/8 dilutions.
 [0076] Determination of the preservation of the concentrated and diluted solution activity, during 2 weeks.
 [0077] Instrument deterioration test.
 [0078] The methodology and results were similar to those of Example 3.
 [0079] Another formula (EXAMPLE 6) with similar results in its 1/1 to 1/6 dilutions in only 5 minutes and 10 minutes
 25 in dilution 1/8, was the following (undiluted formulation):

- hydrogen peroxide: 6.1%
 - lactic acid: 4.8%
 - didecyl-methyl-polyoxy-ethyl-ammonium propionate 50%
 30 - chlorhexidine: 6%
 - silver halides: 0.01%

OTHER EXAMPLES:

- 35 7.1 With other anionic surfactants:

[0080]

7.1.1.:

- 40
 - Hydrogen peroxide: 0.2%
 - Lactic acid: 0.1%
 - Dodecyl sulphate salts. 0.2%
 45 - Ethyl or isopropyl alcohol (60°): until completing 100%

Methodology: it was studied by a contaminated surface model, using a sterile cotton cloth germ-carrier (Her-
 ruzo et al, Rev. Esp. Microbil Clin, 1990; 5:197-202), which was then contaminated with a micro-organism. Then,
 it was introduced in the disinfectant for 5 minutes and removed to submerge it in 5 ml disinfectant inhibitor with
 glass beads. The rest was similar to that described in Example 1 with the endodontic file germ-carrier.

- 50 Results: 4 micro-organisms were used, which are normally found on hospital surfaces: *P. aeruginosa*, *S. au-*
reus, *E. cloacae*, and *C. albicans*, all of them isolated from clinical samples. The results were excellent in only 5
 minutes direct application.

55

Micro-organism	Control (before)	Disinfectant (after application)
<i>P. aeruginosa</i>	2500	0
<i>S. aureus</i>	6900	0

(continued)

Micro-organism	Control (before)	Disinfectant (after application)
E. cloacae	3600	0
C. albicans	4200	0

7.1.2.: The same formula as used in Example 1, but replacing lauryl sulphate by dodecyl sulphate, which in the undiluted formulation is:

- hydrogen peroxide: 16%
- lactic acid: 24%
- dodecyl sulphate salts: 12%
- halogen salts (Br, I): 1%

It was used at 1/1, 1/2 and 1/4 dilutions with Mycobacterium fortuitum 609 and B. subtilis spores (with the bactericidal and sporicidal methodology described in Examples 1 and 2):

Both the concentrated solution and 1/2 and 1/4 dilutions destroyed the inoculii (>5 log10) in 10 minutes.

7.1.3: The same formula as used in Example 1, but replacing lauryl sulphate by alkylsuccinic, which in the undiluted formulation is:

- hydrogen peroxide: 16%
- lactic acid: 24%
- alkyl succinic salts: 12%
- halogen salts (Br, I): 2%

1/1, 1/2 and 1/4 dilutions were also used with Mycobacterium fortuitum and B. subtilis spores (with the bactericidal and sporicidal methodology described in Examples 1 and 2).

Both the concentrated solution and 1/2 and 1/4 dilutions destroyed the inoculii (>5 log 10) in 10 minutes.

7.2. Cationic Surfactants:

[0081]

7.2.1.: A formula was made which in a concentrated solution contained:

- hydrogen peroxide: 22.5%
- lactic acid: 2%
- chlorhexidine: 0.3%
- silver halides 0.0004%

It was also used at 1/1, 1/2 and 1/4 dilutions with Mycobacterium fortuitum 609 and B. subtilis spores (with the bactericidal and sporicidal methodology described in Examples 1 and 2).

The 1/1 and 1/2 concentrated solutions destroyed the inoculii (> 5 log10) in 10 minutes and the 1/4 dilution, destroyed more than 3 log10 in the same time.

7.2.2.: A formula was made, which in a concentrated solution contained:

- hydrogen peroxide: 6.1%
- lactic acid: 4.0%
- ammonium chlorides: 50%
- silver halides: 0.001%

1/1, 1/2, 1/4 and 1/8 dilutions were used with Mycobacterium fortuitum 609 and B. subtilis spores (with the bactericidal and sporicidal methodology described in Examples 1 and 2):

The 1/1 and 1/2 concentrated solutions destroyed the inoculii (> 5 log10) in 5 minutes and the 1/4 and 1/8 dilutions, reached the same effectiveness in 10 minutes.

7.2.3. A formula was made, which in a concentrated solution contained:

- hydrogen peroxide: 6.1%
- lactic acid: 4.0%
- ammonium propyl amide salts: 50%
- silver halides: 0.001%

5 1/1, 1/2, 1/4 and 1/8 dilutions were used with *Mycobacterium fortuitum* 609 and *B. subtilis* spores (with the bactericidal and sporicidal methodology described in Examples 1 and 2):

The 1/1 and 1/2 concentrated solutions destroyed the inoculii (>5 log10) in 5 minutes and the 1/4 and 1/8 dilutions reached the same effectiveness in 10 minutes.

10 In the examples 7.2.2 and 7.2.3, as well as in those described with numbers 2, 3, 4, 5 and 6, the silver salts may be replaced (or added) by iodine, bromine salts or metallic iodine, at a 0.5-5% concentration in the concentrated formulas, with similar results.

On the other hand, any of the previous formulas may be used with enzymes, to obtain enzymatic products, for which we will give some examples, not limiting the possibilities of these formulas:

15 EXAMPLE 8

[0082] The concentrated solution contains:

- hydrogen peroxide: 6.1%
- 20 - lactic acid: 4.5%
- didecyl-methyl-polyoxy-ethyl-ammonium propionate: 20%
- enzymes: trypsin, lipase, amylase: 1-10 ml of each one per ml of final concentrated solution.

25 [0083] It is diluted 100 times and used as a preliminary rinse prior to instrument disinfecting, during 15 minutes. Its effect is checked with the same endodontic file test described previously, against *P. aeruginosa*, *E. cloacae*, *Mycobacterium fortuitum* 609 and commercial *B. subtilis* spores.

[0084] The results are a reduction of more than 2 log10 of all these micro-organisms, while other commercial enzymatic solutions, in which a similar dilution is recommended, only reached 0.1 - 0.9 log 10.

30 EXAMPLE 9

[0085] The concentrated solution contains:

- hydrogen peroxide: 6.1%
- 35 - lactic acid: 4.5%
- lauryl sulphate salts (or dodecyl sulphate): 10%
- enzymes: trypsin, lipase, amylase: 1-10 ml of each one per ml of final concentrated solution.

[0086] The dilutions and results are similar to those of the previous example.

40

PART 2: ANTISEPTICS (prevalent activity)

[0087] Finally, as mentioned in the first page, the composition of this invention may also be used for antiseptics.

45 [0088] The base of this formulation is in the synergy found between a non-chlorinated quaternary ammonium and chlorhexidine and of the latter with halogens (iodine and derivatives). This is shown by means of its MIC (minimum inhibiting concentration).

[0089] A MIC is obtained in a solid medium against *S. aureus* and *P. aeruginosa*.

[0090] The results were the following:

- 50 - Didecyl-methyl-polyoxy-ethyl-ammonium propionate 50% (= Pro)
- Chlorhexidine 20% (= clo)
- Iodine 1% (=I)

55

Product	<i>S. aureus</i>	<i>P. aeruginosa</i>
Pro	1/20480	1/160
Clo	1/20480	1/2560

(continued)

Product	S. aureus	P. aeruginosa
Pro + Clo	1/40960	1/10240
I	1/20	<1/20
Pro + I	>1/40960	1/160
Clo + I	1/40960	1/5120

5 [0091] As may be observed, the MIC increase in the combinations of products indicates synergy between them, since the mixture is diluted to the half with respect to the individual products.

[0092] To determine the bactericidal activity, two classes of tests were used:

- 15 a) "in vitro", following that described by Herruzo et al 1991 (Burns, 1991; 18:35-38).
 b) "in vivo", with the methodology mentioned by Herruzo et al, 2000 (Journal of Surgical Research 2000; 94: 6-12) and a modification of this to artificially contaminate the hands.

Basically, the tests are as follows:

- 20 a) In vitro: 0.5 mm diameter lyophilised pig skin (commercial) circles are cut and sterilised. Afterwards, they are submerged in a micro-organism for 1 hour and left to dry in an inclined position during 15 minutes. Then, they are introduced in the antiseptic during 1 or 3 minutes and finally, the microbicidal is inactivated with the inactivator used in the previous test (6% Tween 80, plus 0.5% sodium bisulphite, plus 0.5% sodium thiosulphate). The rest is like that described in the test with the endodontic files.
- 25 b) In vivo without artificial contamination of the hands: The 4 index-little finger of the non-predominating hand are submerged in 10 ml of the previously mentioned disinfecting inhibitor and shaken inside the inhibitor liquid for 1 minute. Then, cultured in 0.1 ml Mueller-Hinton and extend over the plate. This is the initial natural contamination control. Then, the hand is rinsed with water to eliminate the inhibitor and then rinse with the product to be tested for 3 minutes, rubbing it between both hands (without a surgical brush). Afterwards, the fingers of the predominating hand, etc. are submerged to compare the number of cfu of each one.
- 30 c) In vivo with artificial contamination of the hands: 2 ml of a 24 hour culture of P. aeruginosa are applied over them, letting them to dry for 5 minutes. Afterwards, continue as in the previous experiment, but in the control dilutions are also made before sowing the plate, to be able to make an adequate count of the number of cfu (generally 1000-10000).

35 [0093] The formulas used were (without this being limiting):

EXAMPLE 10:

40 [0094]

- Didecyl-methyl-polyoxy-ethyl-ammonium propionate: 5-20%
- chlorhexidine 1-5%
- iodine or iodinated salts: 0.1-5%
- 45 - emollients: 1-2%
- water, until completing: 100%

EXAMPLE 11:

50 [0095]

- Didecyl-methyl-polyoxy-ethyl-ammonium propionate: 5-20%
- chlorhexidine: 1-5%
- iodine or iodinated salts: 0.1-5%
- 55 - emollients: 1-2%
- alcohol 60°, until completing 100%

EXAMPLE 12: (to be used on persons with allergy to iodine):

[0096]

- 5 - Didecyl-methyl-polyoxy-ethyl-ammonium propionate: 5-20%
- chlorhexidine: 7.5%
- emollients: 1-2%
- water, until completing 100%

10 EXAMPLE 13 (to be used by persons with allergy to iodine):

[0097]

- Didecyl-methyl-polyoxy-ethyl-ammonium propionate: 5-20%
- 15 - chlorhexidine: 7.5%
- emollients: 1-2%
- 60° alcohol, until completing 100%

[0098] The results with respect to 10% iodinated povidone (PVPI), the antiseptic most used in surgical rinsing, were:

20

a) In vitro test:

[0099]

25

Micro-organism	Control Mean cfu	Example, 1 min Mean cfu
P. aeruginosa	1400	No. 10 = 2 No. 11 = 0 No. 12 = 0 No. 13 = 0 PVPI = 75
S. aureus	4600	No. 10 = 0 No. 11 = 0 No. 12 = 0 No. 13 = 0 PVPI = 25

30

35

40

Micro-organism	Control Mean cfu	Example, 3 min Mean cfu
P. aeruginosa	1400	No. 10 = 0 No. 11 = 0 No. 12 = 0 No. 13 = 0 PVPI = 0
S. aureus	4600	No. 10 = 0 No. 11 = 0 No. 12 = 0 No. 13 = 0 PVPI = 12

45

50

55

b) In vivo studies (healthy volunteers n = 15)

[0100]

5

b.1) Normal flora		
	initial (log 10 cfu) (mean E±S)	after rinsing (log 10 cfu) (mean E±S)
	4.24±0.3	Example 10 0 Example 11 <0 Example 12 < 0 Example 13 < 0 PVPI 0.48±0.3

10

15

b.2) Acquired flora (P. aeruginosa)	initial (log 10 cfu) (mean E±S)	after rinsing (log 10 cfu) (mean E±S)
	5.26±0.4	Example 10 0.3±0.3 Example 11 <0 Example 12 0.1±0.1 Example 13 < 0 PVPI 1.98±0.3

20

25

30

35

40

[0101] As may be observed, it is more effective than the reference antiseptic, although the difference is appreciated even more if we study the persistence of the antiseptic effect for two hours under a surgical glove (where there is sweating and recontamination of the hand surfaces as from the micro-organisms housed in the bottom of the sebaceous glands, pillose follicles, etc.). For this purpose, a test is performed, similar to that described in the "in vivo" methods with natural flora, but samples are taken of the natural contamination of both hands, after rinsing and taking samples from the predominating hand. Then, the hands are dried (with the help of a collaborator) with a sterile towel and surgical gloves worn. After two hours, the volunteer removes the glove and a sample taken from the non-predominating hand, in the same way as in the previous trial. Therefore, we have three results: before, immediately after and at 2 hours.

[0102] The results at 2 hours were excellent (no micro-organisms surviving: log 10<0) in examples 11 and 13 (that is, in those with alcoholic solution) and in 10 and 12, the flora obtained increased log 10 with respect to that obtained after rinsing, but still more than 3 log10 below the initial result. However, PVPI, the reference antiseptic, hardly has residual power, since the log10 cfu obtained after removing the glove at 2 hours is 2-3 units greater than that after rinsing and even, sometimes, exceeds that obtained before rinsing, indicating the risk of contamination of the surgical environment in the case of puncturing the glove after this surgical intervention time.

[0103] Not significantly different effects have been obtained by replacing this quaternary ammonium by other non-chlorinated quaternary ammonium salts (data not shown).

Claims

45

50

55

1. A disinfecting and antiseptic composition, **characterised in that** it comprises a mixture of oxidising substances, like hydrogen peroxide, lactic acid plus halogen salts (Br, I) and or heavy metal salts (for example, silver halides), surfactants, whether cationic of the biguanidine type, like chlorhexidine and/or chlorinated or non-chlorinated quaternary ammonium salts or, in some cases, anionic ones (like lauryl sulphate, alkyl succinic or dodecyl sulphate), eventually plus metallic iodine or its salts, together with suitable excipients, some of which may be alcohol.
2. A composition according to claim 1, comprising from 0.2-2.9% or from 6.1-22.5% hydrogen peroxide with respect to the total in its undiluted formulations.
3. A composition according to claim 1, comprising 0.1-24% lactic acid with respect to the total in its undiluted formulations.
4. A composition according to claim 1, comprising 3.1-20% lauryl sulphate salts, with respect to the total in its undiluted formulations.

5. A composition according to claim 1, comprising 0.2-12% dodecyl sulphate salts, with respect to the total in its undiluted formulations.
6. A composition according to claim 1, comprising 0.2-12% alkyl succinic salts, with respect to the total in its undiluted formulations.
7. A composition according to claim 1, comprising 5.1-18% biguanidines like chlorhexidine, with respect to the total in its undiluted formulations.
8. A composition according to claim 1, comprising 0.01-18% biguanidines, like chlorhexidine, with respect to the total in its undiluted formulations.
9. A composition according to claim 1, comprising 0.2-50% quaternary ammonium salts like didecyl-methyl-polyoxy-ethyl-ammonium propionate, ammonium chlorides or compounds of ammonium propylamide, with respect to the total in its undiluted solutions.
10. A composition according to claim 1, comprising 0.2-50% non-chlorinated quaternary ammonium salts like didecyl-methyl-polyoxy-ethyl-ammonium propionate, with respect to the total in its undiluted formulations.
11. A composition according to claim 1, comprising 0.0002-0.1% heavy metal salts, like silver halides, with respect to the total in its undiluted formulations.
12. A composition according to claim 1, comprising 0.1-5% iodine or its salts with respect to the total in its undiluted formulations.
13. A composition according to claim 1, comprising 1-10 units/ml final mixtures of enzymes, like trypsin and/or amylase and/or lipase in its undiluted formulations.
14. A composition according to claim 1, comprising:
 - hydrogen peroxide: 0.2-2.9% or 6.1-22.5%
 - lactic acid: 0.1-4.9% or 15.1-24%
 - lauryl sulphate: 3.1-20%
 - halogen salts (Br, I): 0.1-5%
 - water and excipients (balance until 100%).

where all the percentages are by weight with respect to the total composition in the undiluted formulations.

15. A composition according to claim 1, comprising:
 - hydrogen peroxide: 0.2-2.9% or 6.1-22.5%
 - lactic acid: 0.1-24%
 - dodecyl sulphate or alkyl succinic: 0.2-12%
 - halogen salts (Br, I): 0.1-5%
 - water and excipients (balance until 100%)

where all the percentages are by weight with respect to the total composition in the undiluted formulations.

16. A composition according to claim 1, comprising:
 - hydrogen peroxide: 0.2-2.9% or 6.1-22.5%
 - lactic acid: 0.1-24%
 - chlorhexidine: 5.1-18%
 - heavy metal salts, for example, silver halides: 0.00001-0.1%

where all the percentages are by weight with respect to the total composition in the undiluted formulations.

17. A composition according to claim 1, comprising:

- hydrogen peroxide: 0.2-2.9% or 6.1-22.5%
- lactic acid: 0.1-24%
- didecyl-methyl-polyoxy-ethyl-ammonium propionate: 0.2-50%
- heavy metals salts selected from silver halides: 0.00001-0.1%
- water and excipients (balance until 100%)

where all the percentages are by weight with respect to the total composition in the undiluted formulations.

18. A composition according to claim 1, comprising:

- hydrogen peroxide: 0.2-2.9% or 6.1-22.5%
- lactic acid: 0.1-24%
- didecyl-methyl-polyoxy-ethyl-ammonium propionate: 0.2-50%
- chlorhexidine: 5.1-18%
- heavy metal salts selected from silver halides: 0.00001-0.1%
- water and excipients (balance until 100%)

where all the percentages are by weight with respect to the total composition in the undiluted formulations.

19. A composition according to claim 1, comprising the formulations of claims 15-19 to which enzymes are incorporated, selected from trypsin and/or amylase and/or lipase, to give an enzymatic detergent.

20. A composition according to claim 1, comprising:

- didecyl-methyl-polyoxy-ethyl-ammonium propionate or other non-chlorinated quaternary ammonium salts: 0.2-50%
- chlorhexidine: 0.1-18%
- iodine or its salts: 0.1-5%
- water and excipients (balance until 100%)

where all the percentages are by weight with respect to the total composition in the undiluted formulations.

21. A composition according to claim 1, comprising:

- didecyl-methyl-polyoxy-ethyl-ammonium propionate or other non-chlorinated quaternary ammonium salts: 0.2-50%
- chlorhexidine: 0.1-18%
- iodine or its salts: 0.1-5%
- ethyl or isopropyl alcohol (60°) (balance until 100%)

where all the percentages are by weight with respect to the total composition in the undiluted formulations.

22. A method to disinfect instruments for hospital use, comprising the application of a suitable amount of a disinfecting and antiseptic composition according to any of the claims 1 to 20, on the instruments to be disinfected.

23. A method to disinfect conduits and surfaces of hospital environment, comprising the application of a suitable amount of a disinfecting and antiseptic composition according to any of the claims 1 to 20, on the conduits and surfaces to be disinfected.

24. A method to disinfect equipment, instruments, conduits and surfaces of industrial facilities, comprising the application of a suitable amount of a disinfecting and antiseptic composition according to any of the claims 1 to 20, on the equipment, instruments, conduits and surfaces of the industrial facility to be disinfected.

25. A method and products with or without alcohol, with or without iodine, for skin antiseptics in the human or veterinary field, according to claims 1, 21 and 22.

Figure 1: Assessment of the synergy of "Example a" 1/6 with respect to its basic components (at the same dilution), by means of the determination of the sporicidal effect over a commercial B subtilis germ-carrier





Figure 2 : Assessment of the synergy of "example b" diluted 1/4 with respect to its basic components (at the same dilution), by means of the determination of the sporidical effect over a commercial B subtilis germ-carrier.





European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 02 38 0119 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	HERRUZO CABRERA R.: "Spanish disinfectants for the 21st century" ANALES DE LA REAL ACADEMIA NACIONAL DE MEDICINA, vol. 117, no. 4, 2000, pages 791-812, XP002213667 Spain *Resumen* * page 791 * * page 803: table 2 * *Peroxidina* * page 795 *	1,22-24	A01N37/36 A01N25/30 A61L2/18 A01N59/16 //A01N59/00, 37:36,25:30, 59:16
D, X	EP 1 027 827 A (OFTRAI S L ;MANDALAY INVERSIONES S L (ES)) 16 August 2000 (2000-08-16) * the whole document*	1-4,7,8, 14,22-24	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A01N A61L
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
MUNICH		17 September 2002	Nopper-Jaunky, A
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO FORM 503 (03.02.02) (P04007)

European Patent
OfficeINCOMPLETE SEARCH
SHEET CApplication Number
EP 02 38 0119Claim(s) searched completely:
2-25Claim(s) searched incompletely:
1

Reason for the limitation of the search:

The terms "like", "whether ...or", "in some cases" and "eventually" used in claim 1 have not limiting effects on the scope of protection of said claim so that the features following those terms are to be regarded as entirely optional. Thus, present claim 1 includes every composition comprising a mixture of oxidising substances and surfactants. As a result, a large number of possible compositions are covered by claim 1 and a complete search over the whole of the claimed scope is impossible. Consequently, the search has been carried out only for compositions supported by examples 1-9 of the description, namely compositions comprising at least hydrogen peroxide +lactic acid as oxidising substances. It is to be noted that examples 10 to 13 are not covered by claim 1 since they do not comprise a mixture of oxidising substances (either they just comprise iodine as an oxidising substance or are free of any oxidising substance).



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 02 38 0119

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	EP 0 252 278 A (HENKEL KGAA) 13 January 1988 (1988-01-13) * page 2, line 1 - line 3 * * page 2, line 36 - page 3, line 8; table 1 * * examples 2-4,6,12-16 * * page 8, line 1 - line 11 * ---	1-3,8,25	
X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 94-260811 XP002091567 "Liq bleaching detergent compsn having stability of hydrogen peroxide during storage - contg hydrogen peroxide, surfactants and mixt of glycolic and lactic acid, useful as toilet cleaner. " & JP 06 192692 A (KAO CORPORATION), 12 July 1994 (1994-07-12) * abstract *	1,22-24	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
X	GB 2 076 286 A (QUINODERM LTD) 2 December 1981 (1981-12-02) * claims 1-3,6,8; examples 1,2 *	1-3,25	
X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 1984-267237 XP002213668 "Domestic detergent composition containing anionic and /or nonionic surfactant, organic acid, hydrogen peroxide and water" & JP 59 164400 A (NIPPON PEROXIDE CO LTD), 17 September 1984 (1984-09-17) * abstract *	1,2, 22-24	

	-/--		

EPO FORM 1503 03.82 (P04C10)



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

EP 02 38 0119

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	EP 0 185 970 A (HENKEL KGAA) 2 July 1986 (1986-07-02) * page 1, line 5 - line 9 * * page 2, line 35 - page 3, line 8 * * page 3, line 22 - page 4, line 13 * * claims 1-3,7,8; examples 3,4 * ----	1-25	
A	EP 0 423 922 A (ECOLAB INC) 24 April 1991 (1991-04-24) * page 2, line 4 - line 6 * * page 2, line 51 - page 3, line 4 * * examples 1-9; table 1 * ----	1-25	
A	US 3 297 456 A (NEWELL OWEN W) 10 January 1967 (1967-01-10) * examples 1,2 * ----	1-25	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
A	US 2 917 428 A (HITZMAN DONALD O) 15 December 1959 (1959-12-15) * examples 12,24,36; table 3 * -----	1-25	

EPO FORM 1503 03/82 (P/C10)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 38 0119

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information

17-09-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1027827	A	16-08-2000	WO 9960859 A1	02-12-1999
			AU 7432698 A	13-12-1999
			BR 9815564 A	28-02-2001
			EP 1027827 A1	16-08-2000
			JP 2002516259 T	04-06-2002
			NO 20000385 A	24-02-2000
EP 0252278	A	13-01-1988	DE 3702983 A1	10-12-1987
			AT 169797 T	15-09-1998
			CA 1277899 A1	18-12-1990
			DE 3752210 D1	24-09-1998
			EP 0252278 A2	13-01-1988
			ES 2118705 T3	01-10-1998
			JP 2109426 C	21-11-1996
			JP 8018939 B	28-02-1996
			JP 62292709 A	19-12-1987
			US 4900721 A	13-02-1990
JP 6192692	A	12-07-1994	NONE	
GB 2076286	A	02-12-1981	GB 2076285 A	02-12-1981
			DK 226981 A ,B,	24-11-1981
			US 4485091 A	27-11-1984
JP 59164400	A	17-09-1984	JP 1489788 C	07-04-1989
			JP 63040478 B	11-08-1988
EP 0185970	A	02-07-1986	DE 3445901 A1	19-06-1986
			AT 49695 T	15-02-1990
			BR 8506270 A	26-08-1986
			DE 3575516 D1	01-03-1990
			EP 0185970 A1	02-07-1986
EP 0423922	A	24-04-1991	AT 146036 T	15-12-1996
			AU 623022 B2	30-04-1992
			AU 5783090 A	26-04-1991
			CA 2021631 A1	18-04-1991
			DE 69029360 D1	23-01-1997
			DE 69029360 T2	10-04-1997
			EP 0423922 A1	24-04-1991
			FI 97855 B	29-11-1996
			GB 2236951 A	24-04-1991
			NZ 234345 A	25-02-1992
			US 5139788 A	18-08-1992
US 3297456	A	10-01-1967	NONE	

EPO FORM P0438

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)